

Scar correction with laser

They can hardly be avoided in life: Injuries that leave a more or less noticeable scar after healing. If these scars are unattractively discolored or even significantly thickened, they not only lead to functional limitations, but can also represent a psychological burden.

The fractional CO₂-laser punches tiny holes in the skin, leaving the surrounding tissue intact. As a result, the body produces new collagen and elastin and fills in the skin from below. Scars and hyperpigmentation fade away and the skin appears more even.

Striae (stretch marks)

The fractional laser therapy can be used to improve stretch marks in a particularly gentle and effective way. Stretch marks always appear when the skin is strongly stretched in a relatively short time, for example during pregnancy or weight gain. With the laser beam, the skin is slightly injured and begins a healing process. During this healing, skin-tightening collagen and elastin are formed, which firm the skin in the treated areas and improve elasticity. As a result, the stretch marks fade.

Cesarean scars

Depending on the type of injury, the condition of the wound edges, the wound hygiene, but also depending on age and genetic factors, different visible scars may remain. In most cases, the cesarean section scar heals very nicely, but hypertrophic scars or keloid scars can also occur. Some severe scars require surgical correction. Others can be corrected by laser treatment or with the help of platelet-rich plasma (PRP).

Implementation

Treatment with the latest CO₂ laser technology is painless in most cases, so that even larger areas can be treated without any problems. No prior local anesthetic measures are necessary. The treatment takes between 15 and a maximum of 60 minutes depending on the size and extent.

The cost per session is CHF 300. 1-4 sessions are recommended at intervals of 4 weeks. The treatment is not covered by health insurance.